



## Technical Update 08

### Pea & bean weevil

January 2025

Pea and bean weevil can be a problem in combining, vining or fresh market peas, broad beans, and field beans. Damage caused by the adults is visible as notching around the leaf margins. This damage does not significantly affect yield, but eggs laid during feeding produce larvae which feed in the nitrogen-fixing root nodules of peas and beans, sometimes resulting in yield loss.

#### LIFE CYCLE

The adult weevils are beetle-like in shape and 4 - 5 mm in length. They are light grey to brown in colour with faint striping along the length of the wing cases. They have a very short 'snout' but have the 'elbowed' antennae typical of weevils. Adults migrate from their over-wintering sites, mainly around the grassy uncultivated edges of fields previously cropped with peas or beans. Migration occurs early in the spring, and this often coincides with short periods where the maximum air temperature exceeds 12°C. Both peas and beans may be affected by feeding, although spring sown crops which have recently emerged are more susceptible to damage. Winter beans are usually too far advanced in their growth to be affected by the adults or larvae.



During feeding, eggs laid by the female weevil are washed into the soil around the stem bases and produce larvae which begin feeding inside the root nodules. After 3 - 4 weeks the larvae pupate, and newly emerged adults return to overwintering sites. There is one generation each year.

#### MONITORING FOR WEEVIL

If leaf damage is anticipated, early treatment of the crop is necessary to interrupt the egg laying period and reduce root damage. A monitoring system is available which detects adults when they begin migrating in the early spring. The system was developed by Rothamsted Research and the field development was undertaken by PGRO in conjunction with Rothamsted Research and ADAS. The monitoring system comprises five cone traps containing a pheromone lure. It can be used as an aid to decision making in the following ways:

- Identifies seasons where weevil numbers are low, and crops do not require treatment.
- Identifies the time of peak activity and allows crops to be treated at the optimal time.
- Reduces the need for prophylactic spraying.
- Allows the selection of drilling time to avoid periods of serious damage (useful in organic crop production systems).
- An aid in integrated pest management systems for assured produce crops.

Traps are sited on a single grassy verge or headland of a field which had been cropped with peas or beans the previous year. **They should be sited by mid-February and weevils counted three times each week.** Full instructions are supplied with the traps.

The system is available from Koppert UK <https://www.koppert.com/pea-and-bean-weevil-trap/>.

## CONTROL

### For spring sown peas, field, and broad beans

A **threshold** occurs when the average number of weevils per trap exceeds 30 on a single occasion. Monitoring should continue until a threshold catch is reduced **or** until the latest sown crops have emerged (whichever is the sooner).

When a threshold has been reached, crops which have just emerged or will emerge during the next 10 days may be at risk. A spray can be applied as soon as the first sign of notching is observed and if previous experience is that weevil damage occurs regularly. During periods of slow growth, a second spray should be applied 10 - 14 days later. Crops emerging later should not be at risk. Winter-sown beans seldom require treatment. If a threshold is not reached or if it occurs more than 10 days before crop emergence there is no need to spray. At the end of the season, the lures should be disposed of, and the traps stored for re-use the following year with replacement lures. In recent years pea and bean weevil has developed partial resistance to pyrethroid active substances in some areas.

The information in this publication must not be reproduced without the express written permission from the  
PGRO.

Information disseminated by the Processors & Growers Research Organisation is given after the exercise of all possible care in compilation, preparation and issue, but is provided without liability in its application or use.

## Processors & Growers Research Organisation

The Research Station, Great North Road, Thornhaugh, Cambridgeshire PE8 6HJ, UK  
**Chief Executive:** Roger Vickers **Tel:** +44 (0)1780 782585 **email:** info@pgro.org **www:** pgro.org  
Company Limited by Guarantee - Registered in London No. 567232. Registered Charity No 284077. VAT Reg No 281 7742 37GB